

UNITARY CONTINUOUS REPRESENTATIONS OF COMPACT QUANTUM GROUPS

Bernhard Drabant and Wolfgang Weich

*Max-Planck-Institut für Physik, Föhringer Ring 6, D-80805 München
Sektion Physik der Universität München, Theresienstraße 37, D-80333 München*

Abstract. Generalizing the notion of continuous Hilbert space representations of compact topological groups we define unitary continuous corepresentations of C^* -completions of compact quantum group Hopf algebras on arbitrary Hilbert spaces. It is proved that the unitary continuous corepresentations decompose in finite dimensional irreducible corepresentations.

1 Introduction

Continuous Symmetries play a fundamental rôle in physics. In quantum mechanics the symmetries are represented by unitary operators on the Hilbert space of states which is in general infinite dimensional. This is the physical motivation to study the unitary continuous representations of topological groups. With the quantum groups a more general concept of symmetry has been introduced into physics [1, 2, 6]. They are not any more sets of transformations but are described by non-commutative non-cocommutative $*$ -Hopf algebras which should be understood in some sense as the polynomials over the fictive quantum group. Unitary finite dimensional corepresentations of quantum groups have been studied in several papers, e. g. [4, 5, 7]. The $*$ -Hopf algebras with C^* -norms can be completed to C^* -algebras thus describing non-commutative topological spaces [6] — in the case of commutative $*$ -Hopf algebras the C^* -completions are the algebras of continuous functions over compact topological groups. For an application in quantum mechanics one is now interested in the unitary continuous corepresentations on arbitrary Hilbert spaces of those C^* -algebras arising from non-commutative non-cocommutative $*$ -Hopf algebras.

In section 2 corepresentations of $*$ -Hopf algebras and the theory of $*$ -Hopf algebras of compact quantum groups are reviewed [4, 6]. With the help of the Haar functional on a $*$ -Hopf algebra of a compact quantum group the Hilbert space completion is introduced in section 3. The left action by multiplication yields a C^* -norm. These concepts are used in section 4 to define the continuous corepresentations of the C^* -completion. In the concluding theorem it is stated that

the unitary continuous corepresentations on any Hilbert space decomposes into finite dimensional corepresentations of the underlying $*$ -Hopf algebra. This is a generalization of the classical results on unitary representations of compact topological groups [3].

2 The compact quantum group algebra $\mathcal{A}$

In this section definitions and results of [4, 6] are quoted. Henceforth $\mathcal{A}$ is a $*$ -Hopf algebra with comultiplication $\Delta : \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$, counit $\epsilon : \mathcal{A} \rightarrow \mathcal{C}$ and antipode $S : \mathcal{A} \rightarrow \mathcal{A}$.

For some vector space V let $\text{End}(V)$ denote its endomorphisms. The transposition operator σ_V acts as endomorphism on $V \otimes V$ by $a \otimes b \mapsto b \otimes a$.

2.1 Definition. A corepresentation (T, V) of $\mathcal{A}$ on the vector space V is given by

- (i) $T \in \mathcal{A} \otimes \text{End}(V)$, such that
- (ii) $T_{13}T_{23} = (\Delta \otimes id)(T)$, $T_{23} := 1_{\mathcal{A}} \otimes T$, $T_{13} := (\sigma_{\mathcal{A}} \otimes id)(T_{23})$,
- (iii) $(\epsilon \otimes id)(T) = 1_{\text{End}(V)}$.

If V is a Hilbert space the corepresentation (T, V) of $\mathcal{A}$ is called unitary if $T^*T = 1_{\mathcal{A}} \otimes 1_{\text{End}(V)} = TT^*$, where $T^* := (* \otimes *) (T)$.

If $\dim(V) = n < \infty$ one can choose a basis $\{e_i\}_{i=1, \dots, n}$ and a dual basis $\{e_i^*\}_{i=1, \dots, n}$ with $e_i^*(e_j) = \delta_{ij}$. One can expand $T = \sum_{i,j=1}^n t_{ij} \otimes (e_i \otimes e_j^*)$ and obtains in particular $\Delta(t_{ij}) = \sum_k t_{ik} \otimes t_{kj}$ and $\epsilon(t_{ij}) = \delta_{ij}$. If V is a Hilbert space, (T, V) a unitary corepresentation, then $e_i^* = (e_i | \cdot)$ and $S(t_{ij}) = t_{ji}^*$.

2.2 Definition. Let (T, V) be a corepresentation of $\mathcal{A}$ on the vector space V . A linear subspace W of V is called invariant if $T(1_{\mathcal{A}} \otimes W) \subset \mathcal{A} \otimes W$.

(T, V) is called irreducible if V and $\{0\}$ are the only invariant subspaces of V .

Let (S, W) be another corepresentation of $\mathcal{A}$. A linear operator $L : V \rightarrow W$ is called an intertwiner if $S(1_{\mathcal{A}} \otimes L) = (1_{\mathcal{A}} \otimes L)T$.

(T, V) and (S, W) are called equivalent if there exists a bijective intertwiner.

2.3 Schur's Lemma. Let $L : V \rightarrow W$ be an intertwiner for the corepresentations (T, V) and (S, W) of $\mathcal{A}$. If both are irreducible then either $L = 0$ or L is bijective.

2.4 Definition. The collection of all equivalence classes of irreducible unitary finite dimensional corepresentations of $\mathcal{A}$ is called $\hat{\mathcal{A}}$.

Any $*$ -Hopf algebra has in particular the scalar unitary corepresentation $(1_{\mathcal{A}}, \mathcal{C})$. We denote its equivalence class by $\hat{0} \in \hat{\mathcal{A}}$. For each $\alpha \in \hat{\mathcal{A}}$ we choose a representative (T^α, V^α) which is unitary.

2.5 Proposition. For each $\alpha \in \hat{\mathcal{A}}$ let $\{t_{ij}^\alpha\}_{i,j}$ be the set of matrix elements of (T^α, V^α) . Then $\{t_{ij}^\alpha\}_{\alpha \in \hat{\mathcal{A}}, i, j}$ is a linear independent set.

Setting $\mathcal{A}^\alpha := \text{span}\{t_{ij}^\alpha\}_{i,j} \subset \mathcal{A}$ for $\alpha \in \hat{\mathcal{A}}$ the following definition is motivated by the theory of compact topological groups.

2.6 Definition. A $*$ -Hopf algebra $\mathcal{A}$ is a compact quantum group algebra if it is the linear span of the matrix elements of its unitary irreducible finite-dimensional corepresentations: $\mathcal{A} = \bigoplus_{\alpha \in \hat{\mathcal{A}}} \mathcal{A}^\alpha$.

The algebra of characteristic functions on a compact topological group is such a compact quantum group algebra. The Hopf algebras of the quantum groups $SU_q(N)$ and $SO_q(N)$ ($q > 0$) of [2] are non-commutative examples.

Having a decomposition of $\mathcal{A}$ the Haar functional can be defined.

2.7 Definition. The linear functional $h : \mathcal{A} \rightarrow \mathbb{C}$ with $h(1_{\mathcal{A}}) = 1$ and $h|_{\mathcal{A}^\alpha} = 0$ for $\alpha \neq \hat{0}$ is called the Haar functional.

2.8 Theorem. The Haar functional h on the compact quantum group algebra $\mathcal{A}$ is uniquely determined by its properties

$$(h \otimes id)(\Delta(a)) = h(a) 1_{\mathcal{A}} = (id \otimes h)(\Delta(a)) \quad \text{and} \quad h(1_{\mathcal{A}}) = 1. \quad (1)$$

The Haar functional is positive: $h(a^*a) > 0$ if $a \neq 0$.

3 The completions of $\mathcal{A}$

In sections 3. and 4. $\mathcal{A}$ is always understood to be a compact quantum group algebra with Haar functional h . This Haar functional defines an inner product $\mathcal{A} \times \mathcal{A} \ni (a, b) \mapsto (a|b) := h(a^*b)$ which makes $\mathcal{A}$ a pre-Hilbert space. Thus also $\mathcal{A} \otimes \dots \otimes \mathcal{A}$ becomes naturally a pre-Hilbert space. The Hilbert space completion of the pre-Hilbert spaces X and Y is denoted by $X \hat{\otimes} Y$, the Hilbert space completion of the direct sum of the pre-Hilbert spaces X_i , $i \in I$ is denoted by $\hat{\bigoplus}_{i \in I} X_i$. For a normed vector space V we denote its continuous endomorphisms with $B(V)$.

3.1 Definition. For each $n \in \mathbb{N}$ let $\ell^{\hat{\otimes} n}$ be the Hilbert space completion of the n -fold tensor product of $\mathcal{A}$, let $\ell^{\hat{\otimes} 0} := \mathbb{C}$.

$$\mathbf{L} := \bigoplus_{n \in \mathbb{N}_0}^{\wedge} \ell^{\hat{\otimes} n}. \quad (2)$$

$\ell := \ell^{\hat{\otimes} 1}$ corresponds to the square integrable functions over the quantum group.

$\mathcal{A}$ acts on itself by left multiplication. This yields a $*$ -representation $\mathcal{A} \ni a \mapsto \lambda(a) \in B(\mathcal{A})$, $\lambda(a) : b \mapsto ab \in \mathcal{A}$. Because of the unitarity condition $\sum_i t_{ij}^{\alpha*} t_{ik}^{\alpha} = \delta_{jk}$ in $\mathcal{A}^\alpha$ one gets that

$\lambda(a)$ is a bounded operator on the pre-Hilbert space $\mathcal{A}$ and can therefore be extended to ℓ , i. e. $\mathcal{A} \ni a \mapsto \lambda(a) \in B(\ell)$ is a faithful Hilbert space representation of $\mathcal{A}$ [4]. Cuntt and comultiplication are $*$ -algebra homomorphisms and serve to define Hilbert space representations of $\mathcal{A}$ on $\ell^{\hat{\otimes} n}$ and $\mathbf{L}$.

3.2 Definition. $\Delta^{(0)} := \epsilon$. For $n > 0$, $\Delta^{(n)} := (\Delta^{(n-1)} \otimes id) \circ \Delta$. $\Lambda^{(n)} := \lambda^{\otimes n} \circ \Delta^{(n)} : \mathcal{A} \rightarrow B(\ell^{\hat{\otimes} n})$. $\Lambda := \bigoplus_{n=0}^{\infty} \Lambda^{(n)} : \mathcal{A} \rightarrow B(\mathbf{L})$ is the direct sum Hilbert space representation of $\mathcal{A}$ on $\mathbf{L}$.

This justifies the following definition.

3.3 Definition. The operator norm $|a| := \sup_{x \in \mathbf{L}, \|x\|_{\mathbf{L}}=1} \|\Lambda(a)x\|_{\mathbf{L}} = \|\Lambda(a)\|_{\mathbf{L}}$ is a C^* -norm on $\mathcal{A}$. The corresponding completion of $\mathcal{A}$ is the C^* -algebra $\mathbf{A}$. $\Lambda(\mathbf{A})$ and $\mathbf{A}$ are identified.

3.4 Proposition. Comultiplication and coint on $\mathcal{A}$ are bounded and can be extended to continuous mappings on $\mathbf{A}$, $\Delta : \mathbf{A} \rightarrow \overline{\mathbf{A} \otimes \mathbf{A}}_{\text{closed}} \subset B(\mathbf{L} \otimes \mathbf{L})$, $\epsilon : \mathbf{A} \rightarrow \mathcal{C}$.

Proof. The boundedness of ϵ follows from $|a| = \|\Lambda(a)\|_{\mathbf{L}} \geq \|\Lambda(a)\|_{\ell^{\hat{\otimes} 0}} = |\epsilon(a)|$. The boundedness of Δ comes from $\|(\Lambda \otimes \Lambda)(\Delta(a))\|_{\mathbf{L} \hat{\otimes} \mathbf{L}} = \|\Lambda(a)\|_{\mathbf{L}}$. $\square$

The antipode need not be continuous and can therefore not be extended on $\mathbf{A}$. If $\mathcal{A}$ is the linear span of the matrix elements of the unitary irreducible group representations of a compact topological group then $\mathbf{A}$ is the C^* -algebra of continuous functions over this group. This is the contents of the Peter-Weyl Theorem, see e. g. [3].

With the above construction the compact quantum group algebra $\mathcal{A}$ determines uniquely the C^* -completion $\mathbf{A}$.

3.5 Lemma. The Haar functional h is bounded on $\mathcal{A}$ respective to the norms $\|\cdot\|_{\ell}$ and $|\cdot|$. Its extension on ℓ is given by $\ell \ni x \mapsto (1_{\mathcal{A}}|x)$. Denoting the extension of h on the C^* -algebra by $h : \mathbf{A} \rightarrow \mathcal{C}$ the following holds

$$\forall a \in \mathbf{A} : h(a) = (1_{\mathcal{A}}|\Lambda(a)1_{\mathcal{A}}). \quad (3)$$

4 Continuous corepresentations

Now we can define continuous corepresentations of $\mathbf{A}$. Since $\sigma_{B(L)}$ is bounded on $B(\mathbf{L}) \otimes B(\mathbf{L}) \subset B(\mathbf{L} \hat{\otimes} \mathbf{L})$ it can be continued on the closure.

4.1 Definition. A continuous corepresentation (U, H) of $\mathbf{A}$ on the Hilbert space H is given by

$$(i) \quad U \in \overline{\mathbf{A} \otimes B(H)}_{\text{closed}} \subset B(\mathbf{L} \hat{\otimes} H), \text{ such that}$$

$$(ii) \quad U_{13}U_{23} = (\Delta \otimes id)(U), \quad U_{23} := 1_{\mathbf{A}} \otimes U, \quad U_{13} := (\sigma_{\mathbf{A}} \otimes id)(U_{23}),$$

$$(iii) \quad (\epsilon \otimes id)(U) = 1_{B(H)}.$$

Let U^* denote the adjoint of U in $B(\mathbf{L} \hat{\otimes} H)$.

The continuous corepresentation (U, H) of $\mathbf{A}$ is called isometric if $U^*U = 1_{\mathbf{A}} \otimes 1_{B(H)}$.

(U, H) is called unitary if $UU^* = 1_{\mathbf{A}} \otimes 1_{B(H)} = U^*U$.

In the case of a commutative $*$ -Hopf algebra $\mathcal{A}$ standing for the polynomials over a compact topological group G this definition is equivalent to the statement that the mapping $G \times H \rightarrow H$, $(g, v) \mapsto U(g)(v)$ is a continuous representation of G on H .

If H is finite dimensional then $U \in \mathbf{A} \otimes B(H)$. It is evident that every corepresentation of $\mathcal{A}$ on a Hilbert space is in particular a continuous corepresentation of $\mathbf{A}$. The definition 2.2 of invariant subspaces and irreducibility can be taken over to the case of $\mathbf{A}$.

4.2 Definition. Let (U, H) be a continuous corepresentation of $\mathbf{A}$. The closed subspace $K \subset H$ is called invariant if the orthogonal projector $P_K \in B(H)$, $P_K H = K$ fulfills

$$U(1_{\mathbf{A}} \otimes P_K) = (1_{\mathbf{A}} \otimes P_K)U(1_{\mathbf{A}} \otimes P_K) =: U_K. \quad (4)$$

In this case (U_K, K) is called a continuous subcorepresentation of $\mathbf{A}$.

The continuous corepresentation (U, H) is called irreducible if the only invariant subspaces of H are H and $\{0\}$.

Let (U, H) and (Q, N) be continuous corepresentations of $\mathbf{A}$ on the Hilbert spaces H and N , let $L : H \rightarrow N$ be linear and continuous. If

$$Q(1_{\mathbf{A}} \otimes L) = (1_{\mathbf{A}} \otimes L)U \quad (5)$$

then L is called an intertwiner.

These definitions are justified by the following observation which can easily be verified.

4.3 Lemma. The continuous subcorepresentation (U_K, K) is a continuous corepresentation. If (U, H) is isometric then each continuous subcorepresentation is also isometric.

4.4 Schur's Lemma. Let $L : H \rightarrow N$ be an intertwiner for the corepresentations (U, H) and (Q, N) of $\mathbf{A}$ on the Hilbert spaces H and N with closed image $L(H) \subset N$. If both are irreducible then either $L = 0$ or L is bijective.

Proof. Define in $B(H)$ the orthogonal projector P_0 , $P_0 H = \ker L$ and in $B(K)$ the orthogonal projector P_1 , $P_1 N = L(H)$. Apply to equation (5) $1_{\mathbf{A}} \otimes P_0$ from the right or $1_{\mathbf{A}} \otimes P_1$ from the left respectively to prove that $\ker L$ and $L(H)$ are invariant. If $L \neq 0$ then $\ker L = \{0\}$ because of the irreducibility of (U, H) . If $H \neq \{0\}$ it then follows from the irreducibility of (Q, N) that $N = L(H)$. $\square$

4.5 Lemma. Let H be a finite-dimensional Hilbert space, (U, H) an irreducible isometric continuous corepresentation of $\mathbf{A}$.

- (i) $U \in \mathcal{A} \otimes \mathbf{B}(H)$, and (U, H) is an irreducible corepresentation of $\mathcal{A}$.
- (ii) (U, H) is unitary, $(S \otimes id)(U) = U^*$.

Proof. Let $\{e_A\}_{A=1,\dots,d}$ be an orthonormal basis in H , and $e_A^* := (e_A|\cdot)$ the dual basis. Expand $U = \sum_{A,B=1}^d u_{AB} \otimes (e_A \otimes e_B^*) \in \mathbf{A} \otimes \mathbf{B}(H)$. Consider the following matrices $M_{iA}^{(\alpha,j,B)} := h(t_{ji}^{\alpha*} u_{BA})$. There is at least one $\alpha_u \in \hat{\mathcal{A}}$ and j, B such that $M^{(\alpha_u,j,B)} \neq 0$. The opposite would imply the contradiction

$$\begin{aligned}
0 &= \sum_B \sum_{\alpha,i,j} h(u_{BA}^* t_{ij}^\alpha) \frac{1}{\|t_{ij}^\alpha\|_\ell^2} h(t_{ij}^{\alpha*} u_{BA}) \\
&= \sum_B \sum_{\alpha,i,j} (\Lambda(u_{BA}) 1_A | t_{ij}^\alpha) \frac{1}{\|t_{ij}^\alpha\|_\ell^2} (t_{ij}^\alpha | \Lambda(u_{BA}) 1_A) \\
&= \sum_B (\Lambda(u_{BA}) 1_A | \Lambda(u_{BA}) 1_A) = \sum_B (1_A | \Lambda(u_{BA}^* u_{BA}) 1_A) = 1
\end{aligned}$$

where Lemma 3.5 and the isometry of (U, H) have been used. In V^{α_u} an orthonormal basis $\{f_i\}$ with $f_i^* := (f_i|\cdot)$ such that $T^{\alpha_u} = \sum_{ij} t_{ij}^\alpha \otimes (f_i \otimes f_j^*)$ is used. Then $L := \sum_{iA} M_{iA}^{(\alpha_u,j,B)} (f_i \otimes e_A^*)$ is a non-vanishing intertwiner $(1_{\mathbf{A}} \otimes L)U = T^{\alpha_u}(1_{\mathbf{A}} \otimes L)$. Since $L(H)$ is closed Schur's Lemma implies that L is invertible, and one gets $U = (1_{\mathcal{A}} \otimes L^{-1})T^{\alpha_u}(1_{\mathcal{A}} \otimes L) \in \mathcal{A} \otimes \mathbf{B}(H)$ thus proving the first statement. Applying the antipode to the last equation one gets $(S \otimes id)(U) = (1_{\mathcal{A}} \otimes L^{-1})T^{\alpha_u*}(1_{\mathcal{A}} \otimes L) = U^*$ and hence the unitarity of U . $\square$

Statement (i) of Lemma 4.5 implies that the $*$ -Hopf algebra can be uniquely reconstructed from its C^* -completion $\mathbf{A}$.

4.6 Proposition. Let (U, H) be an isometric continuous corepresentation on the Hilbert space H , K a finite dimensional invariant subspace such that the continuous subcorepresentation (U_K, K) is irreducible.

- (i) (U_K, K) is a unitary continuous corepresentation of $\mathbf{A}$.
- (ii) $U(1_{\mathbf{A}} \otimes P_K) = (1_{\mathbf{A}} \otimes P_K)U$.
- (iii) $H \ominus K$ is invariant, $((1_{\mathbf{A}} \otimes (1_{\mathbf{B}(H)} - P_K))U, H \ominus K)$ is an isometric continuous corepresentation of $\mathbf{A}$.
- (iv) If H is finite dimensional then it decomposes in irreducible continuous subcorepresentations, and $U \in \mathcal{A} \otimes \mathbf{B}(H)$.

Proof. The first statement follows from Lemmas 4.3 and 4.5. One gets therefore $1_{\mathbf{A}} \otimes P_K = U_K U_K^* = U(1_{\mathbf{A}} \otimes P_K)U^*$. Multiplying with U from the right proves the second statement. The third one is a direct consequence using Lemma 4.3. The last assertion follows by induction. $\square$

The last statement means that the finite dimensional unitary continuous corepresentations of $\mathbf{A}$ coincide with the finite dimensional unitary corepresentations of $\mathcal{A}$.

4.7 Definition. Let $f \in \mathcal{A}$, (U, H) an isometric continuous corepresentation of $\mathbf{A}$ on the Hilbert space H .

$$h_f := (h \otimes id)((f \otimes 1_{\mathbf{B}(H)})U) \in \mathbf{B}(H). \quad (6)$$

4.8 Proposition. Let (U, H) be an isometric continuous corepresentation of $\mathbf{A}$ on the Hilbert space H . Let $\{(U_{K_\rho}, K_\rho)\}_{\rho \in R}$ be a family of irreducible finite dimensional continuous subcorepresentations, let $P_\rho := P_{K_\rho}$ be the orthogonal projectors on K_ρ respectively.

- (i) The norm of h_f is bounded by $\|h_f\|_H \leq \|f\|_\ell$.
- (ii) $P_\rho h_f = h_f P_\rho$.
- (iii) $U(1_{\mathbf{A}} \otimes h_f) = \sum_A S(f'_A) \otimes h_{f''_A}$ where $\Delta(f) = \sum_A f'_A \otimes f''_A$.
- (iv) If $v \neq 0$ then there is an $f \in \mathcal{A}$ such that $h_f(v) \neq 0$.

Proof. Apply U to $1_{\mathbf{A}} \otimes v \in \ell \hat{\otimes} H \subset \mathbf{L} \hat{\otimes} H$. $U(1_{\mathbf{A}} \otimes v) =: \sum_i u_i \otimes v_i \in \ell \hat{\otimes} H$ such that the $v_i \in H$ are mutually orthogonal. Then $h_f(v) = \sum_i (f^*|u_i)v_i$. The first assertion follows from $\|h_f(v)\|_H^2 = \sum_i |(f^*|u_i)|^2 \|v_i\|_H^2 \leq \sum_i (f^*|f^*)(u_i|u_i) \|v_i\|_H^2 = \|f^*\|_\ell^2 \|v\|_H^2$. The second is obvious. $U(1_{\mathbf{A}} \otimes h_f) = (id \otimes h \otimes id) \left((1_{\mathbf{A}} \otimes f \otimes 1_{B(H)}) U_{13} U_{23} \right) = (id \otimes h \otimes id) \sum_A \left((\epsilon(f'_A) 1_{\mathbf{A}} \otimes f''_A \otimes 1_{B(H)}) (\Delta \otimes id)(U) \right)$ proves the third statement. $h_f(v) = 0$ is equivalent to $(f^*|u_i) = 0$ for all i . Since $\mathcal{A}$ is dense in ℓ the forth statement follows. $\square$

4.9 Theorem. Let (U, H) be an isometric continuous corepresentation of $\mathbf{A}$ on the Hilbert space H . Then (U, H) is unitary and H decomposes in finite dimensional irreducible continuous subcorepresentations (U_ρ, K_ρ) , $U_\rho \in \mathcal{A} \otimes B(K_\rho)$:

$$H = \bigoplus_{\rho \in R}^\wedge K_\rho. \quad (7)$$

Proof. Using Zorn's Lemma let R be a maximal index set such that P_ρ with $\rho \in R$ are projectors on finite dimensional irreducible subspaces of H with $P_\rho P_{\rho'} = 0$ if $\rho \neq \rho'$. Then choose $v \in (1_{B(H)} - \sum_{\rho \in R} P_\rho)H$. From proposition 4.8 it follows that $h_f(v)$ is contained in a finite dimensional continuous corepresentation. Hence v is necessarily 0. Since the restriction of (U, H) to each irreducible continuous subcorepresentation is unitary the full continuous corepresentation is necessarily unitary. $\square$

The theory of unitary continuous corepresentations of the C^* -completions of the $*$ -Hopf algebras of compact quantum groups is thus reduced to the corepresentation theory of the underlying $*$ -Hopf algebras. In [7] S. L. Woronowicz showed that the theory of finite dimensional corepresentations of the $*$ -Hopf algebra of $SU_q(N)$ coincides with the theory for the undeformed $SU(N)$. Together with our results one concludes therefore that also the theory of unitary continuous corepresentations of these quantum groups coincides with the theory for the undeformed Lie groups.

The authors acknowledge gratefully discussions with M. Dijkhuizen, T. Koornwinder and J. Wess.

References

- [1] V. G. Drinfeld, Proceedings of the International Congress of Mathematicians, Berkeley, California, 798 (1986)
- [2] L. D. Faddeev, N. Yu. Reshetikhin, L. A. Takhtajan, *Algebra and Analysis* **1**, 178 (1989)
- [3] A. W. Knap, Lie groups, Lie algebras, and cohomology, *Mathematical Notes* **34**, Princeton 1988
- [4] T. H. Koornwinder, Lectures at the European School of Group Theory, Trento 1993
M. S. Dijkhuizen, Thesis, CWI Amsterdam 1993
- [5] M. Rosso, *Comm. Math. Phys.* **117**, 581 (1988)
- [6] S. L. Woronowicz, *Comm. Math. Phys.* **111**, 139 (1987)
- [7] S. L. Woronowicz, *Invent. math.* **93**, 35 (1988)